



Environmental influences on healthcare expenditures: an exploratory analysis from Ontario, Canada

M Jerrett, J Eyles, C Dufournaud and S Birch

J. Epidemiol. Community Health 2003;57;334-338
doi:10.1136/jech.57.5.334

Updated information and services can be found at:
<http://jech.bmjournals.com/cgi/content/full/57/5/334>

These include:

References

This article cites 37 articles, 8 of which can be accessed free at:
<http://jech.bmjournals.com/cgi/content/full/57/5/334#BIBL>

Rapid responses

You can respond to this article at:
<http://jech.bmjournals.com/cgi/eletter-submit/57/5/334>

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article

Topic collections

Articles on similar topics can be found in the following collections

[Environmental Issues](#) (621 articles)
[Other Health Policy](#) (547 articles)

Notes

To order reprints of this article go to:
<http://www.bmjournals.com/cgi/reprintform>

To subscribe to *Journal of Epidemiology and Community Health* go to:
<http://www.bmjournals.com/subscriptions/>

PUBLIC HEALTH POLICY AND PRACTICE

Environmental influences on healthcare expenditures: an exploratory analysis from Ontario, Canada

M Jerrett, J Eyles, C Dufournaud, S Birch

J Epidemiol Community Health 2003;57:334–338

See end of article for authors' affiliations

Correspondence to: Professor M Jerrett, School of Geography and Geology, Health Studies Program, and Institute of Environment and Health, McMaster University, Hamilton, Ontario, Canada L8S 4L8; jerrettm@mcmaster.ca

Accepted for publication 26 September 2002

Study objective: This paper explores the relation between healthcare expenditures (HCEs) and environmental variables in Ontario, Canada.

Design: The authors used a sequential two stage regression model to control for variables that may influence HCEs and for the possibility of endogenous relations. The analysis relies on cross sectional ecological data from the 49 counties of Ontario.

Main results: The results show that, after control for other variables that may influence health expenditures, both total toxic pollution output and per capita municipal environmental expenditures have significant associations with health expenditures. Counties with higher pollution output tend to have higher per capita HCEs, while those that spend more on defending environmental quality have lower expenditures on health care.

Conclusions: The implications of our findings are twofold. Firstly, sound investments in public health and environmental protection have external benefits in the form of reduced HCEs. Combined with the other benefits such as recreational values, investments in environmental protection probably yield net social benefits. Secondly, health policy that excludes consideration of environmental quality may eventually result in increased expenditures. These results suggest a need to broaden the cost containment debate to ensure environmental determinants of health receive attention as potential complements to conventional cost control policies.

The purpose of this paper is to explore the relation between healthcare expenditures (HCEs) and environmental variables. Recent public health disasters of environmental origin such as the *Escherichia coli* O157:H7 outbreak in Walkerton, Ontario—which killed seven people and made more than 2000 others ill¹—have rekindled interest in the linkage between environmental quality, government policies, and social costs. The costs of environmental contamination are multifaceted and potentially include increased HCEs.^{2,3} Relations among environmental conditions, government policies to protect public and ecosystem health, and healthcare costs connect to a larger debate on cost containment in the healthcare system.

Concern over cost containment now drives many health reform policies.^{4–6} In Canada, these concerns have led to reforms aimed at containing healthcare costs by various methods: firstly, by imposing expenditure caps on the physician services expenditures within provincial health budgets⁷; secondly, by constraining the incomes of some physicians through individual physician income caps combined with graduated fee penalties; thirdly, by re-evaluating whether certain procedures should be included or removed from the government fee schedule⁸; fourthly, by assessing whether care givers with less training and lower wages can deliver medically necessary procedures, especially in “underserved” areas such as rural communities⁹; and fifthly, by cutting federal transfer payments for health care, which seems to lead to a higher ratio of private to public provision.¹⁰ In the reforms mentioned, the question of whether societal investments in environmental protection lead to reduced health expenditures remains largely unaddressed. In this paper, we examine the question of whether environmental factors contribute to variation in health expenditures.

METHODS

The study area consists of the 49 regions, counties, and districts of Ontario. These are coterminous with the census

divisions used by Statistics Canada and for convenience are called “counties”. Data for the response variable, per capita HCEs, were extracted from public documents.^{11,12} Data for other variables were derived from the 1991 census of Canada,¹³ the Municipal Financial Information System,¹⁴ and the National Pollutant Release Inventory.¹⁵

Overview of the study design

Earlier research indicates that current HCEs might be a function of many variables,¹⁶ including: per capita gross domestic product (GDP),^{17–19} household income,²⁰ demographic dependency ratios, amount of foreign aid, level of urbanisation,^{16,21} ratio of public to private spending on health care,¹⁰ density of physicians,²² and past HCEs. Biophysical environmental variables and those measuring the societal response to environmental degradation are absent from previous analyses, although growing evidence suggests environmental degradation causes some decline in the human population health and in other species on which humans depend for food and resources.^{23–33} The evidence on the environment-health link is strong enough to warrant research that tests the potential influence of these environmental variables on HCEs.² Furthermore, other variables influence population health (for example, educational attainment, economic security, social support), and the potential influence of these variables must be controlled in analyses that test associations between HCEs and environmental variables.

We have attempted to capture the complex system that determines expenditures in a sequential two stage regression analysis. Our modelling controls for the endogenous relation between past HCEs and mortality (a proxy for relative need among populations) and the influence of mortality on current expenditures. We also control other influences on HCEs to prevent confounding in the relation between HCEs and environmental variables.

In Stage 1, we specify the variables that affect population need (to which the medical care system responds) defined as

age adjusted mortality ratios. Other studies show this measure is a good proxy for health care need,³⁴ especially in the Canadian context.³⁵⁻³⁶ We estimate an equation where age standardised mortality ratios are regressed on population health determinants and on past HCEs (in this case, lagged from the previous budget year, 1990–91). This first equation yields predicted standardised mortality ratios that account for past expenditures and current determinants of population health. We argue that the predicted mortality rates are representative of the current health status, having accounted for past variations in HCEs on a local basis.

The predicted values from Stage 1 are entered as an independent variable in a model that uses 1991–92 health care expenditures as the dependent variable in Stage 2. We also include the supply of physicians per county and an indicator variable representing teaching hospitals as likely determinants of current expenditures. Two environmental variables are included in the second stage: total toxic pollution output per county and broadly defined environmental expenditures made by municipal governments. Implicitly we assume an unmeasured morbidity variable intervenes between environmental variables and health expenditures. Given the current state of knowledge on the health effects of environmental degradation, it would prove difficult to predict morbidity because the health effects are often synergistic, subtle, and difficult to quantify. This framework, while imperfect, still allows for a reasonable exploration of the association between environmental factors and HCEs. The specification for the variables is given below. This is followed by a brief commentary on the construction of each variable and the expected direction of the relation.

Model specification and expected relations

Stage 1 uses the following variables:

Y_{it} = 10 year standardised mortality ratios for men, based on data from 1979 to 1988, in the i th county;

X_{1it} = 1990–91 provincial Ministry of Health expenditures per resident in the i th county;

X_{2it} = prevalence of low income, as a percentage of the total population over 15, in the i th county;

X_{3it} = median household income in 1991 Canadian dollars in the i th county;

X_{4it} = educational location quotient for the i th county;

X_{5it} = primary industry employment location quotient for the i th county;

X_{6it} = manufacturing employment location quotient for the i th county; and

e_{it} = residual term for the i th county.

In Stage 1, lagged expenditures (1 April 1990 to 31 March 1991) are used to represent the possible influence of the healthcare system on current health status. Because health expenditures can affect population health and population health needs can affect expenditures, there is an endogenous relation. By incorporating lagged expenditures from the previous year, we estimate the influence of past expenditures on population health needs. The temporal lag ensures the 1990–91 variable is exogenous to the 1991–92 variables, and this serves as an instrumental variable in that these expenditures can affect our dependent variable of interest (that is, 1991–92 expenditures) but cannot be affected by them.³⁸

Socioeconomic conditions in Stage 1 are represented by the incidence of low income relative to the average family, which proxies for inequality; an educational location quotient that measures the proportion of the adult population that exceeds a high school education in each county compared with the provincial adult population proportion; and median household income variable.³⁷⁻³⁹ (See Jerrett *et al.*³⁷⁻³⁹ for detailed derivations of the variables.) Based on the findings of population health research,³⁷⁻⁴⁰⁻⁴² we expect education and household income variables to relate negatively to mortality, while the low income variable should have a positive relation.

Employment structures can influence population health. We have chosen primary industry as an indicator of dangerous and insecure work, and we expect this variable to relate positively to mortality.⁴³ Manufacturing employment represents safer work environments and generally more stable employment. Both anticipation of unemployment and unemployment itself can adversely affect health or health behaviours.⁴⁴⁻⁴⁵ Based on these and other Canadian findings,⁴⁶⁻⁴⁷ we expect manufacturing to relate negatively to mortality.³⁷

Stage 2 includes the following variables:

Y_{it} = 1991–92 provincial health expenditures per person in the i th county;

Y_{it} = predicted male mortality (from Stage 1) for 1991 in the i th county;

X_{1it} = location quotient of general practitioners and family physicians for the i th county;

X_{2it} = an indicator variable representing the presence of a teaching hospital in the i th county;

X_{3it} = municipal environmentally defensive expenditures per resident, in 1991 Canadian dollars, in the i th county; and

X_{4it} = total toxic pollution emissions to all environmental media, in metric tonnes, in the i th county.

e_{it} = residual term in the i th county.

We include two variables to measure possible supply influences on current health expenditures along with predicted mortality to estimate the need for expenditures. A physicians location quotient, measured as the ratio of family physicians per 1000 residents in the county over the comparable rate through the entire province, is used to estimate the relative availability of medical services.³⁷ The county units of analysis leave open the possibility that patients from one county may seek medical services in another county, particularly for specialised treatments. Large teaching hospitals are more likely to have specialists that draw patients from other counties. Such movement toward specialists must be accounted for in the model, and we have done so with the teaching hospital indicator variable. Furthermore, we expect these facilities to retain more highly paid physicians and more expensive capital equipment than other hospitals resulting in a positive relation with expenditures.⁴⁸

We have included two variables as measures of environmental quality and protection. The first measures total pollution emissions and is derived from Environment Canada's National Pollutant Release Inventory.¹⁵ These data estimate total toxic pollution from sources emitting over 10 tonnes per year and were extensively checked for locational and typographical errors.³⁹ The second includes expenditures made by all municipal governments in a county to defend environmental quality. These broadly defined expenditures incorporate sewer and water works, waste management, emergency planning, and recreational facilities, all supplied through the municipal budget.⁴⁹ In Canada, these municipal expenditures account for more than 60% of total environmental expenditures.⁴⁹

We manually selected the model in Stage 1 based on evaluation of the Mallows's Cp statistic, the adjusted r^2 , the standard error of model prediction, and theoretical considerations. The models were fit using ordinary least squares estimation, and variables were transformed to approximate Gaussian normality. In Stage 2, we tested variables together in a multivariate model.

RESULTS

Figure 1 shows a map of the 1991–92 per capita expenditures in the 49 counties of the study area. The map illustrates that considerable variation in HCEs exists throughout the Province, ranging from a low of \$723 in Haliburton county to a high of \$2960 in Frontenac county. Counties with teaching hospitals are in the highest category.




Figure 1 1991 per capita health expenditures.

Table 1 shows the results from both stages of the model. In Stage 1, there are four significant variables: per capita health expenditures 1990–91, education, manufacturing, and primary industry. These variables are associated with 65% of the variation in male mortality. Other variables were tested, but found to be insignificant. To minimise collinearity, we removed insignificant variables from this stage. All significant health determinant variables take the expected sign. The lagged health expenditures coefficient takes a positive sign, which probably would not indicate that past expenditures have contributed to increased population mortality. Instead, it may suggest that some aspects of the medical care system have responded to population need, or that through some

historical accident, the location of those in need coincides with the location of physicians or medical expenditures.

In Stage 2, we used predicted mortality as an independent variable along with a physicians location quotient, an indicator variable for teaching hospitals, and the environmental variables. The predicted mortality variable (representing population need adjusted for past expenditures) shows a positive, significant relation with current expenditures. The teaching hospital variable is highly significant, but the physician variable is insignificant. This variable may be insignificant because of the mortality and hospital variables adequately represent supply of and need for HCEs. The results show that both toxic pollution and municipal defensive expenditures

Table 1 Results of the two stage regression analysis

Variable	Coefficient	SE	t Value	Prob > t	VIF	Partial r
Stage 1 Predicting mortality as a proxy for need controlling for past expenditures						
Constant	205.51	20.42	16.74	0.000		
Health expend t-1	0.011544	0.004556	2.53	0.015	1.1	0.36
Education	-101.24	21.24	-4.77	0.000	1.7	-0.59
Primary industry*	4.72	2.13	2.22	0.032	1.4	0.32
Manufacturing	-17.07	6.27	-2.72	0.009	2.0	-0.39
Adj r ² 65.4%						
Stage 2 Predicting current expenditures						
Constant	2604.25	1295.50	2.01	0.051		
Predicted mortality	10.88	2.52	4.32	0.000	1.1	0.56
Teaching hospital	1092.14	121.82	8.97	0.000	1.6	0.81
Toxic pollution*	45.43	12.87	3.53	0.001	1.2	0.48
Environ expend*	-431.54	204.86	-2.11	0.041	1.2	-0.31
Adj r ² sq 67.1%						

*Transformed with natural log.

Key points

- Previous research on the determinants of healthcare expenditures has excluded environmental variables.
- In Ontario total toxic pollution output relates positively and per capita municipal environmental expenditures relate negatively to healthcare expenditures.
- Sound investments in public health and environmental protection may have external benefits in the form of reduced healthcare expenditures.
- Environmental determinants of health should receive attention as potential complements to conventional health care cost control policies.

have a significant relation with current health expenditures. Total toxic pollution output is positively related to health expenditures, while municipal defensive expenditures are negatively related. All four significant variables account for 67% of the variation in current health expenditures. We tested the unstandardised residuals for spatial autocorrelation with the Moran's I. The results showed a negative coefficient (-0.17) with no significant autocorrelation ($p > 0.1$). Visual inspection of the mapped residuals revealed no unusual patterns. Thus the assumption of independent observations required for efficient and unbiased OLS estimation seems to be fulfilled.

We transformed both environmental variables with the natural log because of a strong positive skew, so the coefficients are interpreted as b/Y_{zmean} . Using this method, the results show that one tonne higher pollution, evaluated at the mean value of expenditures, associates with a 0.03 dollars or 3 cents higher per capita health expenditures. A one dollar increase in defensive expenditures is associated with health expenditures that are 31 cents lower per capita and are not accounted for by the mortality in a population or the presence of teaching hospitals. Interpretation of the coefficient times the range of data shows the magnitude of impact from the highest to the lowest county value. This analysis suggests pollution could exert a \$355 difference in per capita health expenditures, when the lowest pollution level is compared with the highest level (0.03 dollars per tonne \times range of 10800 tonnes = \$355), while health expenditures would be about \$200 lower based on a comparison of the lowest to the highest defensive expenditures ($\$-0.31 \times$ range of \$644 = $\$-200$).

DISCUSSION

Our results show that, after control for other variables that influence health expenditures, both total toxic pollution output and per capita municipal environmental expenditures have significant associations with HCEs. Counties with higher pollution output tend to have higher per capita HCEs, while those that spend more on defending environmental quality have lower HCEs. The implications of this finding are twofold. Firstly, investments in environmental protection are likely to yield external benefits in the form of reduced HCEs. Given other benefits associated with environmental expenditures such as recreation values and maintaining the health of other species, municipal expenditures in environmental protection probably yield net social benefits (that is, expenditures of one dollar yield a benefit greater than one dollar).

Secondly, health policy that excludes considerations of environmental quality may result in increased HCEs, which then "crowd out" other socially beneficial investments. This relation may set in motion a cycle of cumulative consequences whereby environmental degradation leads to higher health expenditures which in turn reduce funds available for environmental protection thus leading to more degradation and so on. In theory, this relation will hold, although in the case of Ontario it is complicated by jurisdictional fragmentation. The provincial government makes the majority of health

expenditures, and municipal governments spend more than other levels of government on environmental protection. Yet, the role of provincial grants in stimulating municipal environmental expenditures⁴⁹ suggests that crowding out may still be important. Furthermore, many instances exist where crowding out by health expenditures will have a direct influence on provincial environmental programmes. This analysis implies that while many other factors must be taken into account in the cost containment debate, exclusion of environmental considerations may lead to social disbenefits.

This study represents an exploratory attempt to quantify the relation between HCEs and environmental variables, and the findings must be tempered with methodological caveats. Firstly, the comparatively small units of analysis used here may lead to possible spillover effects in environmental externalities. Enhanced environmental programmes and expenditures in one jurisdiction would be negated if neighbouring regions pursue environmentally damaging policies. We were unable to address this issue directly, but the lack of spatial autocorrelation in the residuals suggests the variables included in our regression models account for spillovers. If this were not the case, we would expect to see some residual autocorrelation.

Secondly, time lags may influence the relations among environmental investments, environmental quality, and health expenditures. For example, environmental expenditures may not immediately improve environmental quality. Improvements in environmental conditions may take some time to influence health. In turn, improvements in health may not affect demand for health services because of lags in physician, patient, or administrative practices. The direct trade off between environmental and health expenditures as shown in our analysis of the regression coefficients probably overstates our ability to predict these complex relations.

Future research using micro-level data relating specific types of spending (for example, traffic pollution reductions) to individual ill health (for example, asthma) would perhaps better capture the complex environment-health relations explored in this ecological analysis. These refinements seem worth pursuing now that this initial research has established the possibility of a relation between environmental variables and HCEs.

ACKNOWLEDGEMENTS

We thank an anonymous reviewer and the editor for comments that improved the paper. We also thank Pat De Luca and Ric Hamilton for assistance with preparing the map. Moe Watkin of the Ontario Ministry of Health and Long Term Care supplied the health expenditures data and gave useful clarifying comments on its use.

Endnotes

(1) SMR data were extracted from previous studies.³⁷ The long run of mortality data is used to proxy need in 1991. We expect this variable to be representative of the long term mortality pattern and hence adequate for measuring need in 1991. Correlations with female mortality rates are quite high, $r = 0.87$, $p < 0.001$, but male rates were found to correlate more closely with socioeconomic health determinants,³⁷ so we have used these rates for the purpose of generating a predicted value for Stage 2.

(2) Data extracted from Ministry of Health^{11, 12} and based on "Part I" classification that includes operation of hospitals and related facilities, Ontario Health Insurance Plan billings, long term care, mental health services, ambulatory and emergency services, and other primary care services. For the 1991-92 budget year, Part I expenditures accounted for \$15 966 796 256 of \$16 525 215 022 total expenditures or about 96.6% of the total.

(3) Data for the Region and District of Sudbury are combined by the Ministry of Health in expenditures reports. Because we were unable to discern which proportion of the expenditures was attributable to each county, we had to eliminate both cases, leaving 47 counties in the final analysis.

.....

Authors' affiliations

M Jerrett, J Eyles, School of Geography and Geology, Health Studies Program, and Institute of Environment and Health, McMaster University, Hamilton, Ontario, Canada

C Dufournaud, Department of Geography, University of Waterloo, Canada

S Birch, Department of Clinical Epidemiology and Biostatistics and Centre for Health Economics and Policy Analysis, McMaster University

Funding: this research was funded by the McMaster Institute of Environment and Health and the Geographic Information Science Health and Environmental Analysis Laboratory, McMaster University.

Conflicts of interest: none.

REFERENCES

- Health Canada**. Waterborne outbreak of gastroenteritis associated with a contaminated water supply, Walkerton, Ontario, May-June 2000. Ottawa: Canada Communicable Disease Report, 2000;26:170-3.
- Ontario Medical Association**. The illness cost of air pollution: summary of findings. Toronto: Ontario Medical Association, 2000.
- Pearce D**, Turner KR. *Economics of natural resources and the environment*. Baltimore: Johns Hopkins Press, 1991.
- Lomas J**. Devolving authority for health care in Canada's provinces: 4. Emerging issue and prospects. *Can Med Assoc J* 1997;156:817-23.
- Barer ML**, Evans RG, Holt M, et al. It ain't necessarily so: the cost implications of health care reform. *Health Aff* 1994;13:88-99.
- Chernomas R**, Sepelri A. An economist's brief guide to the recent debates on the Canadian health care system. *Int J Health Serv* 1994;24:189-200.
- Hurley J**, Lomas J, Goldsmith U. Physician responses to global physician expenditure budgets in Canada: a common property perspective. *Milbank Q* 1997;75:343-62.
- Barer ML**, Lomas J, Sanmartin C. Re-minding our Ps and Qs: medical cost controls in Canada. *Health Aff* 1996;15:216-34.
- Barer ML**, Stoddart GL. Toward integrated medical resource policies for Canada: 8. geographic distribution of physicians. *Can Med Assoc J* 1992;147:617-23.
- Di Matteo L**. The determinants of the public-private mix in Canadian health care expenditures: 1975-1996. *Health Policy* 2000;52:87-112.
- Ministry of Health**. Information and Planning Branch. *Ontario health expenditures 1990/91: analysis of expenditures by major programs, counties and health districts*. Toronto: Community Information Section of the Information, Planning and Evaluation Branch, 1992.
- Ministry of Health**. Information and Planning Branch. *Ontario health expenditures 1991/92: analysis of expenditures by major programs, counties and health districts*. Toronto: Community Information Section of the Information, Planning and Evaluation Branch, 1993.
- Statistics Canada**. *Data documentation for the 1991 census profile series-part A or B*. Ottawa: Supply and Services Canada, 1991.
- Ministry of Municipal Affairs**. *Municipal financial information 1991*. Ottawa: Queen's Printer for Ontario, 1993.
- Environment Canada**. *National pollutant release inventory data base*. 1995 (available <http://www.doe.ca/pdb/npri.html>).
- Di Matteo L**, Di Matteo R. Evidence on the determinants of Canadian provincial government health expenditures: 1965-1991. *J Health Econ* 1998;17:211-29.
- Newhouse JP**. Medical care expenditures: a cross-national survey. *Journal of Human Resources* 1977;12:115-25.
- Hitris T**, Posnett J. The determinants and effects of health expenditure in developed countries. *J Health Econ* 1992;11:173-81.
- Parkin D**, McGuire A, Yule B. Aggregate health care expenditures and national income: Is health care a luxury good? *J Health Econ* 1987;6:109-27.
- Evans RG**, Stoddart GL. Producing health, consuming health care. *Soc Sci Med* 1990;31:1347-63.
- Hansen P**, King A. The determinants of health care expenditure: a cointegration approach. *J Health Econ* 1996;15:127-37.
- Birch S**. The identification of supplier-inducement in a fixed price system of health care provision: the case of dentistry in the United Kingdom. *J Health Econ* 1988;7:129-50.
- Burnett RT**, Dales RE, Raizenne MB, et al. Effects of low ambient levels of ozone and sulfates on the frequency of respiratory admissions to Ontario hospitals. *Environ Res* 1994;65:172-94.
- Burnett RT**, Dales R, Krewski D, et al. Associations between ambient particulate sulfate and admissions to Ontario hospitals for cardiac and respiratory diseases. *Am J Epidemiol* 1994;142:15-22.
- Geschwind SA**, Stolwijk JAJ, Bracken M, et al. Risk of congenital malformations associated with proximity to hazardous waste sites. *Am J Epidemiol* 1992;135:1197-207.
- Davis DL**, Dinse GE, Hoel DG. Decreasing cardiovascular disease and increasing cancer among whites in the United States from 1973 through 1987: good news and bad news. *JAMA* 1994;271:431-7.
- Patrick DL**, Wickizer TM. Community and health. In: Amick B, Levine S, Tarlov A, et al, eds. *Society and health*. New York: Oxford University Press, 1995:46-92.
- Colborn T**, vom Saal FS, Soto AM. Developmental effects of endocrine disrupting chemicals in wildlife and humans. *Environ Health Perspect* 1993;101:378-83.
- Eyles J**. Environmental health research: setting an agenda by spinning our wheels or climbing the mountain? *Health and Place* 1997;3:1-13.
- Krewski D**, Burnett R, Goldberg MS, et al. *Reanalysis of the Harvard Six-Cities Study and the American Cancer Society Study of Air Pollution and Mortality, Phase II: Sensitivity Analysis*. Cambridge, MA: Health Effects Institute, 2000:295.
- Eyles J**, Elliott SJ. Global environmental change and human health. *Canadian Geographer* 2001;45:99-104.
- Kunzli W**. Assessment of deaths attributable to air pollution: Should we use risk estimates based on time series or cohort studies? *Am J Epidemiol* 2001;153:1050-5.
- Health Canada**. *The health and environment handbook for health professionals*. Ottawa: Minister of Public Works and Government Services Canada, 1998.
- Birch S**, Eyles J, Hurley J, et al. A needs-based approach to resource allocation in health care. *Can J Public Policy* 1993;19:68-85.
- Eyles J**, Birch S. A population needs-based approach to health care resource allocation and planning in Ontario. *Can J Public Health* 1993;84:112-17.
- Eyles J**, Birch S, Chambers S, et al. A needs-based methodology for the allocation of health care resources in Ontario. *Soc Sci Med* 1999;33:489-500.
- Jerrett M**, Eyles J, Cole D. Socioeconomic and environmental covariates of premature mortality in Ontario. *Soc Sci Med* 1998;41:33-49.
- Kennedy P**. *A guide to econometrics*. Cambridge, MA: MIT Press, 1994:411.
- Jerrett M**, Eyles J, Cole D, et al. Environmental equity in Canada: an empirical investigation into the income distribution of pollution in Ontario. *Environ Plann A* 1997;29:1777-800.
- Menchik PL**. Economic status as a determinant of mortality among black and white older men: Does poverty kill? *Popul Stud* 1993;47:427-36.
- Feinstein FS**. The relationship between socioeconomic status and health: a review of the literature. *Milbank Q* 1993;71:279-322.
- Kunst AE**, Mackenbach JP. The size of mortality differences in nine industrialized countries. *Am J Public Health* 1994;84:932-7.
- Marshall KM**. A job to die for. *Perspectives on labour and income*. *BMJ* 1996;311:26-31.
- Ferrie JE**, Shipley MJ, Marmot MG, et al. Health effects of anticipation of job change and non-employment: longitudinal data from the Whitehall study. *BMJ* 1995;311:1264-9.
- Dooley D**, Catalano R, Wilson G. Depression and unemployment: panel findings from the epidemiologic catchment area study. *Am J Community Psychol* 1994;22:745-65.
- Liaw K**, Wort SA, Hayes MB. Intraurban mortality variation and income disparity: a case study of Hamilton-Wentworth region. *Canadian Geographer* 1989;33:131-45.
- D'Arcy C**. Unemployment and health: data and implications. *Can J Public Health* 1986;77:124-31.
- UK Department of Health and Social Services**. *Sharing resources for health in England*. London: HMSO, 1976.
- Jerrett M**. Green costs, red ink: determinants of municipal defensive expenditures. *Professional Geographer* 1999;51:115-34.